Symptoms		Influenza ("Flu")	Cold (Viral upper respiratory infection)	Pertussis
Fever		Usually present and high (102-104°F or 39-40°C); typically lasts 3-4 days	Uncommon	Uncommon, If present, typically low-grade
Headache		Very common	Uncommon	Uncommon
Aches and pains, muscle aches, chest discomfort		Common, often severe	Slight to Moderate	Uncommon
Fatigue and weakness		Moderate to Severe; can last up to 14-21 days	Mild	Mild to Moderate
Extreme exhaustion		Very common early in illness	Extremely Rare	Rare
Stuffy or runny nose		Sometimes	Common	Common early in the disease
Sneezing		Sometime	Common	Common early in the disease
C O U G H	**Character**	Non-productive ("dry") cough is typical; night time cough rare	Hacking cough, often productive; night time cough rare; usually responds to cough medications	Variable character; uncontrollable fits / night time cough is common; generally not responsive to cough medications
	Severity	Moderate	Mild to Moderate	Variable; can be mild in adults and very severe in infants and young children
	Duration	Typically 3-7 days; occasionally to 14 days	Typically 3-7 days	Persistent cough, almost always >1 week, usually 2-6 weeks, sometimes 10+ weeks
	Paroxysms	Uncommon	Rare	Common
Infectious Period		1-2 days before symptom onset to 5-10 days after	Variable; typically 4-7 days after symptom onset; can be longer	From start of runny nose phase (before cough) to 21 days after cough onset*

Síntomas		Influenza ("Gripe")	Resfriado (Infección Viral de las Vías Respiratorias Superiores)	Pertussis(Tos Ferina)
Fiebre		Generalmente presente y alto (102-104°F o 39-40°C); típicamente dura 3-4 días	Poco común	Poco común, si presente, típicamente grado bajo
Dolor de Cabeza		Muy Común	Poco común	Poco común
Dolores y molestias, dolores musculares, molestias en el pecho		Común, muchas veces severo	Leve a Moderado	Poco común
Fatiga y Debilidad		Moderado a severo; puede durar a lo largo de 14-21 días	Leve	Leve a Moderada
Agotamiento Extremo		Muy común al principio de la enfermedad	Extremadamente Raro	Raro
Nariz Tapada o Mocosa		A veces	Común	Común al principio de la enfermedad
Estornudando		A veces	Común	Común al principio de la enfermedad
T O S	**Condición**	Tos no productiva ("seca") es típica; tos nocturna rara	Tos seca, a menudo productiva; tos nocturna rara; generalmente responde a los medicamentos para la tos	Carácter variable; ataques incontrolables / tos nocturna es común; generalmente no responde a los medicamentos para la tos
	Gravedad	Moderada	Leve a Moderada	Variable; puede ser leve en adultos y muy severo en bebés y niños pequeños
	Duración	Típicamente 3-7 días; ocasionalmente a 14 días	Típicamente 3-7 días	Tos persistente, casi siempre> 1 semana, generalmente 2-6 semanas, a veces 10+ semanas
	Paroxismos	Poco común	Raro	Común
Periodo infeccioso		1-2 días antes del inicio de los síntomas a 5-10 días después	Variable; típicamente 4-7 días después del inicio de los síntomas; puede ser más largo	Desde el comienzo de la fase nasal líquida (antes de la tos) hasta los 21 días posteriores a la aparición de la tos *